


Amendment to the Claims


The claimed invention is:

1. (Currently Amended) A compound of formula (Ia), (Ib), or (Ic):



or a pharmaceutically acceptable salt, ~~prodrug~~, tautomer, hydrate or ~~solvate~~ thereof, wherein:

R¹ is a group of the formula



wherein R¹ can optionally be further independently substituted with at least one moiety independently selected from the group consisting of: carbonyl, halo, halo(C₁-C₆)alkyl, perhalo(C₁-C₆)alkyl, perhalo(C₁-C₆)alkoxy, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, hydroxy, oxo, mercapto, (C₁-C₆)alkylthio, (C₁-C₆)alkoxy, (C₅-C₁₀)aryl or (C₅-C₁₀)heteroaryl, (C₅-C₁₀)aryloxy or (C₅-C₁₀)heteroaryloxy, (C₅-C₁₀)ar(C₁-C₆)alkyl or (C₅-C₁₀)heteroar(C₁-C₆)alkyl,

(C₅-C₁₀)ar(C₁-C₆)alkoxy or (C₅-C₁₀)heteroar(C₁-C₆)alkoxy, HO-(C=O)-, ester, amido, ether, amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkylamino(C₁-C₆)alkyl, di(C₁-C₆)alkylamino(C₁-C₆)alkyl, (C₅-C₁₀)heterocyclyl(C₁-C₆)alkyl, (C₁-C₆)alkyl- and di(C₁-C₆)alkylamino, cyano, nitro, carbamoyl, (C₁-C₆)alkylcarbonyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₅-C₁₀)arylcabonyl, (C₅-C₁₀)aryloxycarbonyl, (C₁-C₆)alkylsulfonyl, and (C₅-C₁₀)arylsulfonyl;

each R³ is independently selected from the group consisting of: hydrogen, halo, halo(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, perhalo(C₁-C₆)alkyl, phenyl, (C₃-C₁₀)cycloalkyl, hydroxy, (C₁-C₆)alkoxy, perhalo(C₁-C₆)alkoxy, phenoxy, (C₃-C₁₀)cycloalkyl-O-, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-NH-SO₂-, O₂N-, NC-, amino, Ph(CH₂)₁₋₆HN-, (C₁-C₆)alkyl HN-, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂-amino, (C₁-C₆)alkyl-SO₂-NH-, amino(C=O)-, aminoO₂S-, (C₁-C₆)alkyl-(C=O)-NH-, (C₁-C₆)alkyl-(C=O)-[((C₁-C₆)alkyl)-N]-, phenyl-(C=O)-NH-, phenyl-(C=O)-[((C₁-C₆)alkyl)-N]-, (C₁-C₆)alkyl-(C=O)-, phenyl-(C=O)-, (C₃-C₁₀)cycloalkyl-(C=O)-, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, H₂N(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂-N-(C=O)-, phenyl-NH-(C=O)-, phenyl-[(C₁-C₆)alkyl)-N]-(C=O)-, (C₃-C₁₀)cycloalkyl-NH-(C=O)- and (C₁-C₆)alkyl-(C=O)-O-;

where alkyl, alkenyl, alkynyl, phenyl, cycloalkyl, alkoxy, phenoxy, amino of R³ is optionally substituted by at least one substituent independently selected from (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, halo, H₂N-, Ph(CH₂)₁₋₆HN-, and (C₁-C₆)alkylHN-;

s is an integer from one to five;

and

R⁶ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, phenyl, (C₃-C₁₀)cycloalkyl, (C₁-C₆)alkyl-(SO₂)-, phenyl-(SO₂)-, H₂N-(SO₂)-, (C₁-C₆)alkyl-NH-(SO₂)-, ((C₁-C₆)alkyl)₂N-(SO₂)-, phenyl-NH-(SO₂)-, (phenyl)₂N-(SO₂)-, (C₁-C₆)alkyl-(C=O)-, phenyl-(C=O)-, (C₃-C₁₀)cycloalkyl-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, (C₃-C₁₀)cycloalkyl-O-(C=O)-, H₂N-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-, phenyl-NH-(C=O)-, (C₃-C₁₀)cycloalkyl-NH-(C=O)-, ((C₁-C₆)alkyl)₂N-(C=O)-, (phenyl)₂N-(C=O)-, phenyl-[(C₁-C₆)alkyl)-N]-(C=O)-,

and (C₃-C₁₀)cycloalkyl-(((C₁-C₆)alkyl)-N)-(C=O)-;

where alkyl, alkenyl, alkynyl, phenyl, benzyl, cycloalkyl, alkoxy, phenoxy, amino of R⁶ is optionally substituted with at least one moiety independently selected from the group consisting of halo, (C₁-C₆)alkyl,

(C₂-C₆)alkenyl, (C₂-C₆)alkynyl, perhalo(C₁-C₆)alkyl, (C₃-C₁₀)cycloalkyl, phenyl, benzyl, (C₅-C₁₀)heterocyclic, (C₅-C₁₀)heteroaryl, (C₁-C₆)alkyl-SO₂-, formyl, NC-,

(C₁-C₆)alkyl-(C=O)-, (C₃-C₁₀)cycloalkyl-(C=O)-, phenyl-(C=O)-,

(C₅-C₁₀)heterocyclic-(C=O)-, (C₅-C₁₀)heteroaryl-(C=O)-, HO-(C=O)-,

(C₁-C₆)alkyl-O-(C=O)-, (C₃-C₁₀)cycloalkyl-O-(C=O)-,

(C₅-C₁₀)heterocyclic-O-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-,

(C₃-C₁₀)cycloalkyl-NH-(C=O)-, phenyl-NH-(C=O)-,

(C₅-C₁₀)heterocyclic-NH-(C=O)-, (C₅-C₁₀)heteroaryl-NH-(C=O)-,

((C₁-C₆)alkyl)₂-N-(C=O)-, phenyl-(((C₁-C₆)alkyl)-N)-(C=O)-, hydroxy,

(C₁-C₆)alkoxy, perhalo(C₁-C₆)alkoxy, (C₃-C₁₀)cycloalkyl-O-, phenoxy,

(C₅-C₁₀)heterocyclic-O-, (C₅-C₁₀)heteroaryl-O-, (C₁-C₆)alkyl-(C=O)-O-,

(C₃-C₁₀)cycloalkyl-(C=O)-O-, phenyl-(C=O)-O-, (C₅-C₁₀)heterocyclic-(C=O)-O-,

(C₅-C₁₀)heteroaryl-(C=O)-O-, O₂N-, amino, (C₁-C₆)alkylamino,

((C₁-C₆)alkyl)₂-amino, formamidyl, (C₁-C₆)alkyl-(C=O)-NH-,

(C₃-C₁₀)cycloalkyl-(C=O)-NH-, phenyl-(C=O)-NH-,

(C₅-C₁₀)heterocyclic-(C=O)-NH-, (C₅-C₁₀)heteroaryl-(C=O)-NH-,

(C₁-C₆)alkyl-(C=O)-[(((C₁-C₆)alkyl)-N]-, phenyl-(C=O)-[(C₁-C₆)alkyl-N]-,

(C₁-C₆)alkyl-SO₂NH-, (C₃-C₁₀)cycloalkyl-SO₂NH-, phenyl-SO₂NH-,

(C₅-C₁₀)heterocyclic-SO₂NH- and (C₅-C₁₀)heteroaryl-SO₂NH-;

wherein the phenyl moiety of a R⁶ substituent is optionally further substituted with at least one radical independently selected from the group consisting of halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, perfluoro(C₁-C₆)alkyl and perfluoro(C₁-C₆)alkoxy.

2. (Previously Cancelled)

3. (Previously Cancelled)

4. (Previously Cancelled)

5. (Previously Cancelled)

6. (Previously Cancelled)

7. (Previously Cancelled)
8. (Previously Cancelled)
9. (Original) A compound of claim 1, wherein s is one to two; R³ is hydrogen or (C₁-C₆)alkyl; and R⁶ is H, (C₁-C₆)alkyl, or (C₃-C₁₀)cycloalkyl.
10. (Original) A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.
11. (Cancelled)
12. (Cancelled)
13. (Previously Presented) A compound 6-[5-(6-methyl-pyridin-2-yl)-2H-[1,2,3] triazol-4-yl]-quinazoline or a pharmaceutically acceptable salt thereof.
14. (Previously Presented) A pharmaceutical composition comprising 6-[5-(6-methyl-pyridin-2-yl)-2H-[1,2,3] triazol-4-yl]-quinazoline or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.